



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A01N 33/12, 33/08, 25/30 // (A01N 33/08, 61:00, 59:24, 59:00, 25:30) (A01N 33/12, 61:00, 59:24, 59:00, 25:30)	A1	(11) International Publication Number: WO 94/23574 (43) International Publication Date: 27 October 1994 (27.10.94)
(21) International Application Number: PCT/EP94/01180 (22) International Filing Date: 14 April 1994 (14.04.94) (30) Priority Data: 93201138.0 20 April 1993 (20.04.93) EP (34) Countries for which the regional or international application was filed: NL et al. (71) Applicant (for all designated States except US): AKZO NOBEL N.V. [NL/NL]; Velperweg 76, NL-6824 BM Arnhem (NL). (72) Inventors; and (73) Inventors/Applicants (for US only): NORTH, Michael, Shaun [ZA/ZA]; 47 Alkin Street, Brackenfell, 7560 Cape Town (ZA). BUTSELAAR, Robert, Jan [NL/NL]; Zwinglilaan 40, NL-1216 ME Hilversum (NL). (74) Agent: SCHALKWIJK, Pieter, Cornelis; Akzo Nobel N.V., Patent Department (Dept. APTA), P.O. Box 9300, NL-6800 SB Arnhem (NL).		(81) Designated States: AU, BR, NZ, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>

(54) Title: ACTIVITY PROMOTING ADDITIVES FOR REST-BREAKING AGENTS**(57) Abstract**

Activity promoting additives for rest-breaking agents are disclosed. Further, restbreaking compositions comprising these additives and a rest-breaking agent are also disclosed. Finally, a process for breaking the rest of deciduous fruit trees is disclosed. The activity promoting additives enhance the activity of rest-breaking agents thereby leading to improvements in the yields and quality of fruit from deciduous fruit trees in regions which have mild winter weather conditions.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
RJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

Activity Promoting Additives for Rest-Breaking Agents

The present invention relates to the use of particular compounds as activity promoting additives for rest-breaking agents, to rest-breaking compositions comprising these additives and to the use of these compositions for rest-breaking of deciduous fruit trees.

BACKGROUND OF THE INVENTION

Deciduous fruit trees require winter chilling to grow normally. The amount of chilling required depends upon the kind of fruit and the cultivar. If winter chilling is insufficient, then growth abnormalities such as delayed and uneven blossoming, poor leaf cover, insufficient fruit-set and reduced fruit size can occur. These symptoms are generally referred to as delayed foliation according to the article, Blommaert, K.L.J., "Winter Dormancy and Delayed Foliation," The Deciduous Fruit Grower, (1956).

Measures to reduce the symptoms of delayed foliation include treatment with high volumes of chemical rest-breaking agents during later winter and various physical manipulations such as pruning.

In South Africa, for example, most apple trees receive insufficient winter chilling to break rest completely and thus annual application of a chemical rest-breaking agent is standard practice. The most widely used rest-breaking agent in commercial apple orchards is dinitro-ortho-cresol (DNOC) in combination with winter-oil emulsion (a lipophilic agent) (Erez, A. et al., "Improved Methods of Breaking Rest in the Peach and Other Deciduous Fruit Species," J. Amer. Soc. Hort. Sci., 96, pp. 519-522 (1971)). This article also mentions the use of the potassium salt of gibberellic acid, kinetin, indoleacetic acid and thiourea as potential rest-breaking agents.

DNOC is a non-systemic stomach poison and contact insecticide, ovicidal to the eggs of certain insects. It is strongly phytotoxic and its permissible use as an insecticide is limited to dormant sprays or on waste ground. Further, DNOC is known to act as a cumulative poison in man. Although DNOC is relatively inexpensive and effective as a rest-breaking agent, it is toxic enough to limit its continued use and it is currently on the European Red List of agricultural chemicals which will be prohibited in the future. DNOC is also banned in the United States for use as an agricultural chemical.

One of the most effective alternatives to DNOC is hydrogen cyanamide according to the publications, Snir, I., "Chemical Dormancy Breaking of Red Raspberry," HortScience, 18, pp. 719-713 (1983); North, M.S., "Effects of Cyanamide and DNOC/oil on Budbreak, Yield and Fruit Size of Golden Delicious Apples," S. Afr. J. Plant Soil, 6(3), pp. 176-178 (1989); Stadler, J.D., North, M.S. and Lutze, G.F.A., "Artificial Rest-Breaking of Apricot and Plum Cultivars Using Hydrogen Cyanamide," J.S.Afr.Soc.Hort.Sci., 1(1), pp. 9-11, (1991); Nee, C.C. and Fuchigami, L.H., "Overcoming Rest at Different Growth with Hydrogen Cyanamide," Scientia Horticulturae, 50, pp. 107-113 (1992); and North, M.S., "Alternative Rest-Breaking Agents to DNOC/oil for Apples," S. Afr. J. Plant Soil, 9(1), pp. 39-40 (1992).

Hydrogen cyanamide is a skin and eye irritant and is especially acute when used in combination with the consumption of alcohol. The toxicity and relatively high price limit its market acceptance and hydrogen cyanamide has also been placed on the European Red List.

Thus, there exists an immediate need for new, milder rest-breaking agents, not only in South Africa, but also in countries which are currently developing deciduous fruit industries but do not have an ideal climate therefor, such as Brazil. Further, there is a need for

such new rest-breaking agents in countries with existing deciduous fruit industries that, up to now, did not appreciate the extent of the problem of delayed foliation.

5 The response of a tree to a rest-breaking agent is dependent upon the chemical composition of the agent, its application rate and timing and on the nutritional status of the tree according to Terblanche, J.H. and Strydom, D.K., "Effects of Autumnal Nitrogen Nutrition, Urea Sprays and a Winter Rest-Breaking Spray on Budbreak and Blossoming of
10 Young 'Golden Delicious' Trees Grown in Sand Culture," Deciduous Fruit Grower, 23 pp. 8-14 (1973).


Many chemicals have been found to have rest-breaking ability. A
15 summary of some of these chemicals can be found in Saure, M.C., "Dormancy Release in Deciduous Fruit Trees," Janick, J. (Ed.), Horticultural Reviews, 7, pp. 239-287, AVI Publishing Co. Inc., Westport, Connecticut (1985). The efficacy of these various
20 rest-breaking agents appears to be directly related to their physiological harshness.

One of the milder rest-breaking agents is potassium nitrate which has been shown to have a positive effect on peaches. Generally, fruit trees having a lower chill requirement, such as peaches, also require
25 lower concentrations of rest-breaking agents, whereas fruits with a higher chill requirement, such as apples, require excessive concentrations of a mild rest-breaking agent. Articles demonstrating the effects of thiourea, potassium nitrate and combinations thereof include, Wolak, R.J. and Couvillon, G.A., "Time of Thiourea-KNO₃
30 Application on the Rest Requirement and Bud Development in 'Loring' Peach," HortScience, 11(4), pp. 400-402, (1976) and Fernandez-Escobar, R. and Martin, R., "Chemical Treatments for Breaking Rest in Peach in Relation to Accumulated Chilling," J. Hort. Sci., 62(4), pp. 457-461, (1987).


For the foregoing reasons, there exists a need in the art for improved rest-breaking agents which are, affordable, effective, less toxic than the present rest-breaking agents and which can be employed in reasonable concentrations.

Summary of the Invention

The present invention relates to a process for breaking the rest in deciduous fruit trees which comprises the step of applying to at least one deciduous fruit tree before blossom, an effective amount of at least one rest-breaking agent and a compound selected from the group consisting of alkoxylated amines represented by the following general formula:




wherein n is an integer from 1 to 50, A represents an alkylene group and when n>1, each A may be the same or different alkylene groups, R is selected from straight or branched chain alkyl or alkenyl groups having 8 to 22 carbon atoms and groups represented by the formula:




wherein m is an integer from 1-50, A₁ represents an alkylene group and when n>1, each A₁ may be the same or different alkylene groups, x is an integer from 1-6, and R₂ is independently selected from the same groups as R, R₃ is selected from hydrogen, 1-8 carbon atom straight or branched chain alkyl and alkenyl groups and aryl groups having up to 8


carbon atoms; and R_1 is selected from hydrogen, straight or branched chain alkyl and alkenyl groups having 1-22 carbon atoms, a group represented by the formula:



wherein n' is an integer from 1 to 50 and A_2 represents an alkylene group and when $n' > 1$ each A_2 may be the same or different alkylene groups; and alkoxyated quaternary ammonium compounds represented by the following general formula:



wherein n , A , R_1 and R_3 are as defined above, X is an anion, R_5 is selected from hydrogen, straight or branched chain alkyl or alkenyl groups having 1 to 4 carbon atoms, benzyl or R_5X^- can be carboxymethyl as in betaines and oxygen as in amine oxides; R_4 is selected from straight or branched chain alkyl and alkenyl groups having 8-22 carbon atoms and groups represented by the formula:



wherein A_1 , m , R_2 , R_3 , R_4 , R_5 and x are as defined above; to enhance the rest-breaking of the fruit tree.

More particularly, X may be halides such as Cl^- , Br^- , CH_3SO_4^- , and $\text{C}_2\text{H}_5\text{SO}_4^-$, among others. The anion associated with these quaternary ammonium compounds is not critical to the process of the present invention.

These compounds are known from European patent application publication number 0 463 241 where it is disclosed that these compounds are useful as blossom thinning agents for stone fruits. In addition, a small group of the foregoing compounds are known from European patent 0 257 686 which discloses several alkoxylated amines and their use as activity promoting additives for herbicides and fungicides. Also disclosed therein is a method for making these compounds, which method is hereby incorporated by reference.

As a result of extensive research it has now been found that the 5 above-identified compounds can be used as adjuvants for chemical rest-breaking agents for deciduous fruits. These compounds exhibit a substantial effect when combined with rest-breaking agents, and human toxicity and phytotoxicity studies have shown that these compounds 20 have acceptably low levels of toxicity to humans and other plants. Further, the compounds do not cause significant harm to useful insect populations. Finally, the activity-enhancing effect of these compounds allows the use of milder rest-breaking agents which are less toxic, as well as the use of lower concentrations of rest-breaking agents. 25

The amino compounds of the present invention may be prepared by reacting an amine selected from the group consisting of R-NH_2 , RRNH , and $\text{R-NH-((CH}_2)_x\text{-NH)}_n\text{-R'}$ wherein R and R' are aliphatic hydrocarbon groups having 8-22 carbon atoms, $n=1-5$ and x is an integer from 1-6; 30 with at least one alkylene oxide.

The preferred alkylene oxides for use in the present invention are ethylene oxide, propylene oxide, isobutylene oxide and butylene oxide. The compounds of the present invention are made in such a way as to introduce varying numbers of alkylene oxide units onto the amino nitrogen. Thus, these alkylene oxide groups may be all the same, such as, for example, one or more ethylene oxide units, or the groups may be different to form, for example, block copolymer chains of ethylene oxide and propylene oxide units, random copolymer chains consisting of several units of each of two or more different alkylene oxides, or alternating units of two or more alkylene oxides. Any conceivable combination of alkylene oxide units up to 50 units long may be employed at each location on the amino nitrogen which is to contain such units. In addition, a single amino nitrogen may contain two different alkylene oxide chains attached thereto or two chains which are the same.

In the most preferred embodiments of the present invention, block copolymer chains of ethylene oxide and one or more of propylene oxide or butylene oxide are employed. Preferably, the molar weight of the compounds used in the present invention is less than 8000 though higher molecular weight compounds can be employed in some circumstances.

The amino compounds can also be quaternized by known quaternization methods to produce quaternary ammonium compounds which are also useful in the process of the present invention. The fourth substituent added to the amino nitrogen by quaternization may be an alkyl, aryl or alkenyl group having 1 to 4 carbon atoms. The anion associated with such quaternary ammonium compounds is not critical to the process of the present invention.

Typical compounds suitable for use in the process of the present invention include, but are not limited to, cocobis (2-hydroxyethyl)methylammonium chloride, polyoxyethylene (15) cocomethylammonium chloride, oleylbis (2-hydroxyethyl)methylammonium chloride, polyoxyethylene (15) stearyl methylammonium chloride, cocobis (2-hydroxyethyl)amine, polyoxyethylene(5)cocoamine, polyoxyethylene(15)cocoamine, tallowbis (2-hydroxyethyl)amine, polyoxyethylene(5)tallowamine, polyoxyethylene(15)tallowamine, tallow/oleylbis(2-hydroxyethyl)amine, oleylbis(2-hydroxyethyl)amine, polyoxyethylene(5)oleylamine, polyoxyethylene(15)oleylamine, hydrogenated tallowbis(2-hydroxyethyl)amine, hydrogenated polyoxyethylene(5)tallowamine, hydrogenated polyoxyethylene(15)tallowamine, hydrogenated polyoxyethylene(50)tallowamine, N,N',N'-tris(2-hydroxyethyl)-N-tallow-1,3-diaminopropane, N,N',N'-polyoxyethylene(10)-N-tallow-1,3-diaminopropane, and N,N',N'-polyoxyethylene(15)-N-tallow-1,3-diaminopropane.

The process of the present invention is useful for rest-breaking of deciduous fruit to produce improvements advancing the time of bloom, budbreak and/or leaf cover and fruit set. The present process also enables the use of either milder rest-breaking agents or smaller quantities of harsher rest-breaking agents thus providing considerable advantages in toxicity to man and insect populations as well as safety advantages in the application of these agents.

When reference is made in this patent application to enhancement of rest-breaking, what is meant is that the time of bloom is advanced or one or more budbreak and/or leaf cover or fruit set are improved over a control tree.

5 The process of the present invention is to be applied to deciduous fruit trees prior to blossom. The process is particularly useful in growing areas where the winter chilling of the fruit trees is insufficient to provide good budbreak and fruit set for the particular fruit or cultivar. Winter chilling can be measured and is often represented by the unit, "Richardson Chill Units."

10 The optimum time to break rest for a particular deciduous fruit tree will depend upon several factors including the type of fruit, the cultivar, the climatic conditions and the type and amount of rest-breaking agent being applied. For some fruit or cultivar species, the best rest-breaking effects are accomplished by early application of the rest-breaking agent to the trees whereas for others it is best to wait until just before blossom. In general, the rest-breaking agent will be applied at some point between the time when winter has peaked and the time when blossoming begins.

20 One of the effects of the rest-breaking agents of the present invention is to accelerate the blossoming of the fruit trees. Accordingly, in the application of these rest-breaking agents consideration should be given to this effect and the agents should not be applied too early such that blooming occurs when there is still a risk of harsh weather.

25 The process of the present invention is carried out by the application of an effective amount of at least one rest-breaking agent in combination with an amount at least one compound as defined herein to enhance the rest-breaking activity of said rest-breaking agent. The rest-breaking compositions are preferably applied in the form of an aqueous solution in a concentration of 0.25 to 30% and more preferably from 0.5 to 10%. The lower limit is generally determined by the upper limit on application volume for the particular application equipment

30

being employed, as well as by the type of fruit, the cultivar and the particular rest-breaking composition.

5 The upper concentration limit will generally be dictated by phytotoxicity considerations as higher concentrations of certain compounds have a localized phytotoxic effect on the trees. Thus, a concentration should be selected which provides adequate rest-breaking without unwanted phytotoxic effects on the remainder of the tree. The
10 attached examples show that such concentrations can be selected by routine experimentation with the particular species of deciduous fruit.

15 The activity promoting additive of the present invention is generally employed in a concentration of 0.1-10.0% in the aqueous solution and more preferably, 0.2-5.0%. Again, the amount of activity promoting additive required will depend on the fruit, the cultivar and the particular rest-breaking agent, as well as the quantity of rest-breaking agent to be employed. Generally, the less rest-breaking
20 agent employed, the more activity-promoting additive required to maintain the desired level of rest-breaking activity.

Rest-breaking agents which may be used in the process and composition of the invention include any known rest-breaking agent. Examples
25 include hydrogen cyanamide, potassium nitrate, the potassium salt of gibberellic acid, kinetin, indoleacetic acid, thiourea and combinations thereof. The most preferred rest-breaking agent for use in the present invention is potassium nitrate.

30 The composition is preferably applied in a manner similar to the manner in which commercial insecticides are applied. More particularly, conventional equipment such as knapsack sprayers, hand held spray guns, mist blowers, and aerial spraying equipment among

others may be used. The composition is applied the same way as in pesticide application.

5 The process of the present invention has the significant advantages that it breaks rest to the extent that the use of known, highly toxic rest-breaking agents can be eliminated or considerably reduced, it can be done in a manner which is safe for the crops and the treatment has no long term phytotoxic effect on the orchards, if carried out
10 correctly. Further, the rest-breaking process will cause significantly less harm to beneficial insects when applied within the normal application volume, and the process appears to be environmentally acceptable, non-hazardous to operators of the application equipment, and non-corrosive to the equipment.

15 The present invention will be further illustrated by the examples appended hereto.


Examples 1-9 and Comparative Examples A-O

20 Effect of Armoblen® ACER 89002 (ACER) and Winter Oil (Oil) on Hydrogen Cyanamide (HC) Rest-Breaking Agent in Golden Delicious Apples

25 In Examples 1-9, various combinations of Dormex™ (hydrogen cyanamide rest breaking agent) and Armoblen® ACER 89002 (activity promoting additive) in aqueous spray solutions were employed as rest-breaking agents in Golden Delicious apples. These tests were carried out in a region having a winter chill factor of less than 300 Richardson chill units.

30 The treatments of all the field trials in the examples have all been applied by motor driven piston pump equipment fitted with hand held spray lances. Operating pressure was kept at a constant two Bars.

Armoblen® ACER 89002 can be represented by the general formula:



wherein $p=12$, $q=4$ and R is a hydrocarbon group derived from tallow fatty acid.

Comparative Example A is the control example where no rest-breaking agent was applied. Comparative Examples B-D demonstrate the effects of various amounts of hydrogen cyanamide alone. Comparative Examples E-G demonstrate the effects of various amounts of Armoblen® ACER 89002 alone. Comparative Examples H-I demonstrate the effect of various amounts of winter oil when used alone. Comparative Examples J-O demonstrate the effects of various combinations of winter oil and hydrogen cyanamide.

As far as timing of the applications is concerned, all materials were applied at the same time. The amounts of winter oil appear to be high relative to the amount of Armoblen® ACER 89002. However, the amounts of winter oil employed in the examples, namely 3.0-5.0%, are the amounts which are currently employed in practice.

Results:

The following Table I gives the results obtained in these examples. All amounts are volume percent, based on the total volume of the aqueous solution.

The results for budbreak represent the absolute amount of budbreak (amount of budding per tree). The results for fruit set represent the number of fruit per flower cluster on the tree.

TABLE 1

Example	HC	Oil	ACER	Bud Break	Fruit Set
A	0	0	0	16.34	2.39
1	1.0	0	0.1	17.03	1.21
2	2.0	0	0.1	26.28	1.33
3	4.0	0	0.1	25.92	0.80
4	1.0	0	0.5	19.91	2.04
5	2.0	0	0.5	19.38	1.59
6	4.0	0	0.5	26.80	1.34
7	1.0	0	1.0	22.00	1.47
8	2.0	0	1.0	17.24	1.11
9	4.0	0	1.0	20.04	1.09
B	1.0	0	0	19.56	1.13
C	2.0	0	0	22.38	1.15
D	4.0	0	0	27.53	1.13
E	0	0	0.1	15.11	1.65
F	0	0	0.5	21.40	2.34
G	0	0	1.0	13.06	1.83
H	0	3.0	0	14.75	1.60
I	0	5.0	0	18.87	1.65
J	1.0	3.0	0	13.61	1.40
K	2.0	3.0	0	14.19	1.16
L	4.0	3.0	0	12.78	0.97
M	1.0	5.0	0	16.54	1.18
N	2.0	5.0	0	14.47	0.77
O	4.0	5.0	0	9.56	1.49

These results demonstrate that the formulations of the present invention significantly improve the bud break while giving a fruit set of at least the same order as the prior art compositions.

Examples 10-17 and Comparative Examples P-Y

5 Eight year old Golden Delicious trees grown at 1000 trees per hectare were treated with three liters of the following aqueous solutions per tree. The experiment had a completely randomized design and 5 repetitions per treatment were employed. These tests were carried out in a region with a somewhat harsher winter chilling of about 600 Richardson chilling units.

10 In Examples 10-13 and Comparative Examples P-Y, the rest-breaking agents were applied at the same time. In Examples 14-17, the rest-breaking agent KNO_3 was applied 10 days after the application of a solution of only the activity promoting additive ACER.

15 Comparative Example P was the control example. Comparative Examples Q and R were solutions of Armoblen® ACER-89002. Comparative Examples S-T were winter oil emulsions. Comparative Example U was a commercially used formulation of DNOC with winter oil emulsion. Comparative Examples V-Y were commercially used formulations of winter oil and hydrogen cyanamide. Examples 10-17 were Armoblen® ACER-89002 in various combinations with the milder rest-breaking agent potassium nitrate. The results are given in Table 2.

Results

25 For these examples budbreak is expressed as a percentage and fruit set is expressed as the number of fruit per tree.

TABLE 2

	Example	RBA	Amount RBA	Oil	ACER	Days Bloom Advanced	Budbreak (%)	Fruit Set
5	P	---	0	0	0	---	49	504.8
	Q	---	0	0	1.0	5	49	399.4
	R	---	0	0	2.0	10	52	525.2
	S	---	0	2.0	0	10	61.5	283.8
	T	---	0	3.0	0	16	49.5	323.8
	U	DNOC	5.0	80.0	0	16	38	360.4
	V	HC	0.5	2.0	0	22	59.5	531.6
	W	HC	0.5	3.0	0	22	63	457.2
10	X	HC	0.75	2.0	0	27	77.5	669.8
	Y	HC	0.75	3.0	0	27	67	491.8
	10	KNO ₃	4.0	0	1.0	33	45.5	532
	11	KNO ₃	4.0	0	2.0	36	47.5	547
	12	KNO ₃	8.0	0	1.0	22	37.5	717.6
	13	KNO ₃	8.0	0	2.0	36	58.5	1276.2
	14	KNO ₃	4.0	0	1.0	5	44.5	401.4
15	15	KNO ₃	4.0	0	2.0	22	32.5	469
	16	KNO ₃	8.0	0	1.0	16	40	387.6
	17	KNO ₃	8.0	0	2.0	22	42	529.6

These results show that the compositions of the present invention advanced the time of bloom more than that of the compositions of hydrogen cyanamide/winter oil and were significantly better than the DNOC/oil. The activity promoting additive alone (ACER) seemed to delay the time of bloom relative to control trees.

The winter oil and winter oil and cyanamide gave the best results for budbreak. However, the effect of the milder, less toxic rest-breaking agents of the invention is apparent from these tests and shows positive results.

With respect to fruit set, the rest-breaking agents of the invention performed generally better than the harsher commercially applied rest breaking agents and, particularly in Example 13, an exceptional result was obtained.

Examples 18-26 and Comparative Examples Z-MM

5 Five year old Golden Delicious trees planted in high density were treated with three liters of the following aqueous solutions per tree. The experiment had a completely randomized design and 3 repetitions per treatment were employed. These tests were carried out in a region with only mild winter chilling of less than 300 Richardson chilling units.

10 In Examples 18-26 and Comparative Examples Z-CC and MM, the rest-breaking agents were applied at the same time. In Examples DD-LL, the rest-breaking agent KNO_3 was applied 10 days later.

15 Comparative Example Z was the control example. Comparative Examples AA-CC were solutions of KNO_3 . Comparative Examples DD-LL formulations of winter oil and KNO_3 . Examples 18-26 were Armoblen® ACER-89002 in various combinations with KNO_3 . Comparative Example MM was 4% hydrogen cyanamide. The results are given in Table 3.

20

Results

25 For these examples leaf cover is expressed as a rating of from 1-5 with 5 being the best and 1 being the worst. Fruit thinned is expressed as both the grams and number of fruit thinned.

30

TABLE 3

	Example	KNO ₃	Oil	ACER	Days Before Full Bloom	Leaf Cover	Grams Fruit Thinned	No Fruit Thinned-
5	Z	0	0	0	1.66	3.67	51.5	17
	AA	4.0	0	0	6.66	4.00	30.1	8
	BB	8.0	0	0	5.00	3.00	49.7	12
	CC	12.0	0	0	8.33	3.67	69.9	12
	DD	4.0	2.0	0	3.33	3.33	45.5	9
	EE	4.0	4.0	0	3.33	3.00	9.0	3
	FF	4.0	6.0	0	1.66	2.00	0	0
10	GG	8.0	2.0	0	6.66	3.00	25.9	11
	HH	8.0	4.0	0	5.00	3.00	22.6	6
	II	8.0	6.0	0	5.66	2.67	51.6	15
	JJ	12.0	2.0	0	5.00	3.33	49.7	13
	KK	12.0	4.0	0	6.66	2.67	118.6	21
	LL	12.0	6.0	0	6.66	4.00	29.6	4
	MM	*	0	0	18.33	4.00	66.2	8
15	18	4.0	0	0.5	3.33	2.67	12.2	3
	19	4.0	0	1.0	8.33	4.00	77.2	19
	20	4.0	0	2.0	6.66	3.67	149.5	41
	21	8.0	0	0.5	1.66	2.67	43	18
	22	8.0	0	1.0	5.00	3.00	88.2	18
	23	8.0	0	2.0	6.66	3.33	44.6	12
	24	12.0	0	0.5	3.33	2.67	31.1	7
	25	12.0	0	1.0	3.33	3.00	0	0
20	26	12.0	0	2.0	6.66	3.00	43.5	10

* = 4.0% Hydrogen Cyanamide

These results show that the relatively toxic hydrogen cyanamide gives the best results with respect to days before full bloom. However, a treatment of 1% ACER with 4% KNO₃ is shown to be equal in effect to a treatment with 6% winter oil and 12% KNO₃. Thus, the activity promoting additive of the present invention can be used in smaller quantities than the winter oil and allows a significant reduction in the amount of mild rest-breaking agent while increasing the level of rest-breaking activity.

The best results for leaf cover were obtained with the hydrogen cyanamide, 1% ACER plus 4% KNO₃ and 2% winter oil with 12% KNO₃ again

demonstrating that the activity promoting additive of the present invention allows a reduction in the quantity of rest-breaking agent required to achieve a good result.

5 The best results for fruit set were obtained with the 12% KNO_3 , the 1% and 2% ACER with 4% KNO_3 and the 4% winter oil with 12% KNO_3 . Again a significant reduction in the amount of rest-breaking agent is made possible by the present invention.

10 Examples 27-30 and Comparative Examples NN-QQ

15 Nine year old Golden Delicious trees grown at 1000 trees per hectare were treated with three liters of the following aqueous solutions per tree. The experiment had a completely randomized design and 5 repetitions per treatment were employed. These tests were carried out in a region with a somewhat harsher winter chilling of about 600 Richardson chilling units.

20 In all Examples, the rest-breaking agents were applied at the same time. Comparative Example NN was the control example. Comparative Example OO was DNOC/oil and Comparative Examples PP-QQ were ACER control examples. The results are given in Table 4.

25 Results

For these examples budbreak is expressed as a percentage and fruit set is expressed as the absolute number of fruit per tree.

TABLE 4

Example	DNOC/Oil	KNO ₃	ACER	Days Bloom Advanced	Budbreak (%)	Fruit Set Number
5						
NN	0	0	0	----	49	450
OO	5.0	0	0	15.0	39	350
PP	0	0	1.0	4.0	49	400
QQ	0	0	2.0	10.0	52	550
27	0	4.0	1.0	33.0	45	550
28	0	4.0	2.0	37.0	48	620
29	0	8.0	1.0	20.0	43	550
10	30	0	8.0	33.0	53	1200

These results show that the compositions of the present invention were better than the control and DNOC for budbreak and the time that bloom was advanced. Further, the fruit set was also improved by the treatments in accordance with the present invention and in Example 30 a dramatic improvement is observed.

Examples 31-39 and Comparative Examples RR-VV

Six year old Golden Delicious trees planted in high density were treated with three liters of the following aqueous solutions per tree. The experiment had a completely randomized design and 3 repetitions per treatment were employed. These tests were carried out in a region with only mild winter chilling of less than 300 Richardson chilling units.

In all Examples the rest-breaking agents were applied at the same time. Comparative Example RR was the control example. Comparative Example VV was 2.0% hydrogen cyanamide. Comparative Examples SS-TT were ACER control examples. The results are given in Table 5.

Results

For these examples leaf cover is expressed as a rating of from 1-5 with 5 being the best and 1 being the worst. Fruit thinned is expressed as the number of fruit.

TABLE 5


	Example	HC	KNO ₃	ACER	Leafcover	Fruit Set Number
10	RR	0	0	0	3.00	8
	SS	0	4.0	0	3.67	6
	TT	0	8.0	0	2.67	5
	UU	0	12.0	0	3.33	11
	VV	2.0	0	0	4.00	4
	31	0	4.0	0.5	2.67	2
	32	0	8.0	0.5	2.67	7
15	33	0	12.0	0.5	2.67	8
	34	0	4.0	1.0	4.00	15
	35	0	8.0	1.0	3.00	9
	36	0	12.0	1.0	3.00	2
	37	0	4.0	2.0	3.67	22
	38	0	8.0	2.0	3.33	5
	39	0	12.0	2.0	3.00	6

These tests show that the compositions of the present invention give an improvement over the control for leaf cover (an expression representing budbreak) and are as effective as the more toxic hydrogen cyanamide rest-breaking agent for leaf cover (budbreak). Again, the compositions of the present invention gave the best fruit set at 4% KNO₃.


The foregoing examples have been presented for purposes of illustration and description only and are not to be construed as limiting the scope of the invention in any manner. Accordingly, the scope of the invention is to be determined by the claims appended hereto.

What is claimed is:


1. A process for breaking the rest in deciduous fruit trees which comprises the step of applying to at least one deciduous fruit tree before blossom, an effective amount of at least one rest-breaking agent and a compound selected from alkoxyated amines represented by the following general formula:



wherein n is an integer from 1 to 50, A represents an alkylene group and when n>1, each A may be the same or different alkylene groups, R is selected from straight or branched chain alkyl or alkenyl groups having 8 to 22 carbon atoms and groups represented by the formula:




wherein m is an integer from 1-50, A₁ represents an alkylene group and when n>1, each A₁ may be the same or different alkylene groups, x is an integer from 1-6, and R₂ is independently selected from the same groups as R, R₃ is selected from hydrogen, 1-8 carbon atom straight or branched chain alkyl and alkenyl groups and aryl groups having up to 8 carbon atoms; and R₁ is selected from hydrogen, straight or branched chain alkyl and alkenyl groups having 1-22 carbon atoms, a group represented by the formula:



wherein n' is an integer from 1 to 50 and A_2 represents an alkylene group and when $n' > 1$ each A_2 may be the same or different alkylene groups; and alkoxyated quaternary ammonium compounds represented by the following general formula:


5



10

wherein n , A , R_1 and R_3 are as defined above, X is an anion, R_5 is selected from hydrogen, straight or branched chain alkyl or alkenyl groups having 1 to 4 carbon atoms, benzyl or R_5X^- can be carboxymethyl as in betaines and oxygen as in amine oxides; R_4 is selected from straight or branched chain alkyl and alkenyl groups having 8-22 carbon atoms and groups represented by the formula:

15



20

wherein A_1 , m , R_2 , R_3 , R_4 , R_5 and x are as defined above; to enhance the rest-breaking of the deciduous fruit tree.


25

- 2 A process in accordance with claim 1 wherein said rest-breaking agent is applied in the form of an aqueous solution having a concentration of from 0.1 to 10% of said compound and from 0.5-30% of said rest-breaking agent.


30

3. A process in accordance with claim 2 wherein said compound has a molecular weight of less than 8000 grams/mole.


4. A process in accordance with claim 3 wherein said compound is an alkoxyated quaternary ammonium compound represented by the general formula:



wherein n is an integer from 1 to 50, X^- is an anion, A represents an alkylene group and when $n > 1$, each A may be the same or different alkylene groups, R_5 is selected from hydrogen, straight or branched chain alkyl and alkenyl groups having 1-4 carbon atoms, benzyl, or $R_5 X^-$ can be carboxymethyl as in betaines, and oxygen as in amine oxides; R_4 is selected from straight or branched chain alkyl or alkenyl groups having 8 to 22 carbon atoms or a group represented by the formula:




wherein m is an integer from 1 to 50, X^- is an anion, R_5 is as defined above, A_1 represents an alkylene group and when $n > 1$, each A_1 may be the same or different alkylene groups, x is an integer from 1 to 6, R_2 is selected from straight or branched chain alkyl or alkenyl groups having 8-22 carbon atoms; and R_1 is selected from hydrogen, straight or branched chain alkyl or alkenyl groups having 1 to 22 carbon atoms, a group represented by the formula:



wherein n' is an integer from 1 to 50 and A_2 represents an alkylene group; when $n' > 1$ each A_2 may be the same or different alkylene groups, and R_3 is selected from hydrogen, straight or branched chain alkyl or alkenyl groups having 1 to 8 carbon atoms and aryl groups having up to 8 carbon atoms.


5. A process in accordance with claim 3 wherein said compound is an alkoxyated amine represented by the following general formula:



wherein n and n' are integers from 1 to 50, A represents an alkylene group and when n or n' are greater than 1, each A may be the same or different alkylene groups, and R is selected from straight or branched chain alkyl or alkenyl groups having 8 to 22 carbon atoms.


6. A process in accordance with claim 5 wherein $n > 1$ and A includes both ethoxy groups and propoxy groups.

7. A process in accordance with claim 3 wherein said compound is an alkoxyated amine represented by the following general formula:



wherein n and n' are integers from 1 to 50, A represents an alkylene group and when n or n' are greater than 1, each A may be


the same or different alkylene groups, R is a group represented by the formula:



wherein m is an integer from 1 to 50, A₁ represents an alkylene group and when m>1, each A₁ may be the same or different alkylene groups; x is an integer from 1 to 6, and R₂ is selected from straight or branched chain alkyl or alkenyl groups having 8-22 carbon atoms.


8. A process in accordance with claim 3 wherein R is selected from alkyl groups having 12-22 carbon atoms.

9. A process in accordance with claim 3 wherein said compound is an amine oxide represented by the following general formula:




wherein R, A, n and m are as defined in claim 1.


10. Use of a compound represented by the following general formula:




wherein n is an integer from 1 to 50, A represents an alkylene group and when n>1, each A may be the same or different alkylene groups, R is selected from straight or branched chain alkyl or alkenyl groups having 8 to 22 carbon atoms and groups represented by the formula:




wherein m is an integer from 1-50, A₁ represents an alkylene group and when n>1, each A₁ may be the same or different alkylene groups, x is an integer from 1-6, and R₂ is independently selected from the same groups as R, R₃ is selected from hydrogen, 1-8 carbon atom straight or branched chain alkyl and alkenyl groups and aryl groups having up to 8 carbon atoms; and R₁ is selected from hydrogen, straight or branched chain alkyl and alkenyl groups having 1-22 carbon atoms, a group represented by the formula:



wherein n' is an integer from 1 to 50 and A₂ represents an alkylene group and when n'>1 each A₂ may be the same or different alkylene groups; and alkoxylated quaternary ammonium compounds represented by the following general formula:




wherein n, A, R₁ and R₃ are as defined above, X is an anion, R₅ is selected from hydrogen, straight or branched chain alkyl or alkenyl groups having 1 to 4 carbon atoms, benzyl or R₅X- van be carboxy-methyl as in betaines and oxygen as in amine oxides; R₄ is selected from straight or branched chain alkyl and alkenyl groups having 8-22 carbon atoms and groups represented by the formula:



wherein A₁, m, R₂, R₃, R₄, R₅ and x are as defined above; as an activity promoting additive to enhance the activity of a rest-breaking agent.


11. Use in accordance with claim 10 wherein said compound has a molecular weight of less than 8000 grams/mole and is an alkoxyated amine represented by the following general formula:




wherein n and n' are integers from 1 to 50, A represents an alkylene group and when n or n' are greater than 1, each A may be the same or different alkylene groups, and R is selected from straight or branched chain alkyl or alkenyl groups having 8 to 22 carbon atoms.

12. A composition useful for breaking the rest in deciduous fruit trees which comprises an effective amount of at least one rest-


breaking agent and an amount of a compound selected from
alkkoxyated amines represented by the following general formula
which is effective to enhance the activity of said rest breaking
agent:



wherein n is an integer from 1 to 50, A represents an alkylene
group and when n>1, each A may be the same or different alkylene
groups, R is selected from straight or branched chain alkyl or
alkenyl groups having 8 to 22 carbon atoms and groups represented
by the formula:




wherein m is an integer from 1-50, A₁ represents an alkylene group
and when n>1, each A₁ may be the same or different alkylene
groups, x is an integer from 1-6, and R₂ is independently selected
from the same groups as R, R₃ is selected from hydrogen, 1-8
carbon atom straight or branched chain alkyl and alkenyl groups
and aryl groups having up to 8 carbon atoms; and R₁ is selected
from hydrogen, straight or branched chain alkyl and alkenyl groups
having 1-22 carbon atoms, a group represented by the formula:




wherein n' is an integer from 1 to 50 and A₂ represents an
alkylene group and when n'>1 each A₂ may be the same or different

alkylene groups; and alkoxyated quaternary ammonium compounds represented by the following general formula:




wherein n, A, R₁ and R₃ are as defined above, X is an anion, R₅ is selected from hydrogen, straight or branched chain alkyl or alkenyl groups having 1 to 4 carbon atoms, benzyl or R₅X- can be carboxy-methyl as in betaines and oxygen as in amine oxides; R₄ is selected from straight or branched chain alkyl and alkenyl groups having 8-22 carbon atoms and groups represented by the formula:



wherein A₁, m, R₂, R₃, R₄, R₅ and x are as defined above.

13. A composition in accordance with claim 12 wherein said compound has a molecular weight of less than 8000 grams/mole and is an alkoxyated amine represented by the following general formula:



wherein n and n' are integers from 1 to 50, A represents an alkylene group and when n or n' are greater than 1, each A may be

the same or different alkylene groups, and R is selected from straight or branched chain alkyl or alkenyl groups having 8 to 22 carbon atoms.

5

10

15

20

25

30

INTERNATIONAL SEARCH REPORT

 Int. Application No
 PCT/94/01180

 A. CLASSIFICATION OF SUBJECT MATTER
 IPC 5 A01N33/12 A01N33/08 A01N25/30 //(A01N33/08,61:00,59:24,
 59:00,25:30), (A01N33/12,61:00,59:24,59:00,25:30)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

 Minimum documentation searched (classification system followed by classification symbols)
 IPC 5 A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP,A,0 257 686 (AKZO) 2 March 1988 cited in the application see the whole document ---	1-13
Y	DE,A,31 50 404 (SKW TROSTBERG) 30 June 1983 see page 6 - page 7 see claim 3 ---	1-13
Y	GB,A,1 604 321 (R.C.BARBA) 9 December 1981 see the whole document --- -/--	1-13

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

26 July 1994

Date of mailing of the international search report

03.08.94

Name and mailing address of the ISA

 European Patent Office, P.O. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tlx. 31 651 epo nl,
 Fax (+31-70) 340-3016

Authorized officer

Lamers, W

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 94/01180

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	GB,A,2 257 044 (IMPERIAL CHEMICAL INDUSTRIES) 6 January 1993 see page 1, paragraph 3 see page 1, paragraph 5 see page 1, line 24 see page 11; table IV see claims 1,2,21 ----	1-13
X	US,A,4 556 410 (P.M. RONNING ET AL.) 3 December 1985 see column 1, line 6 - column 4, line 48 ----	12,13
X	US,A,4 525 200 (M.KIMPARA ET AL.) 25 June 1985 see column 1, line 6 - column 3, line 3 ----	12,13
X	EP,A,0 272 542 (BASF) 29 June 1988 see page 3, line 1 - page 4, line 20 ----	12,13
A	EP,A,0 463 241 (AKZO) 2 January 1992 cited in the application see the whole document ----	1-13
A	EP,A,0 232 755 (MITSUBISHI GAS CHEMICAL) 19 August 1987 -----	1-13

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 94/01180

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0257686	02-03-88	CA-A- 1329538	17-05-94
		US-A- 5226943	13-07-93
		ZA-A- 8706184	22-02-88
DE-A-3150404	30-06-83	JP-B- 2062524	26-12-90
		JP-A- 58110511	01-07-83
		US-A- 4487625	11-12-84
GB-A-1604321	09-12-81	NONE	
GB-A-2257044	06-01-93	NONE	
US-A-4556410	03-12-85	NONE	
US-A-4525200	25-06-85	JP-B- 4072801	19-11-92
		JP-A- 59157002	06-09-84
EP-A-0272542	29-06-88	DE-A- 3643246	30-06-88
		AU-B- 600059	02-08-90
		AU-A- 8264887	23-06-88
		DE-A- 3784658	15-04-93
		ES-T- 2044903	16-01-94
		US-A- 4936899	26-06-90
EP-A-0463241	02-01-92	AU-B- 635836	01-04-93
		AU-A- 7630991	02-01-92
		CA-A- 2045352	27-12-91
		US-A- 5118340	02-06-92
EP-A-0232755	19-08-87	JP-A- 62167704	24-07-87
		US-A- 4799950	24-01-89

THIS PAGE BLANK (USPTO)